New Jersey Surgery Center PRE-OPERATIVE INTERVIEW

NAME:	M or F	Height	Weight	Age
Procedure to be done				
Emergency contact person and number				
Person who will provide your transportation	n home			
Phone number where we can reach you bet	tween 7am – 3	3pm		
Please list allergies (with reactions) to food	d and/or medi	cations or state	e NONE:	
Do you have latex allergy? Y/N If yes	s, please call u	us at 609-581-6	5230	
PAST MEDICAL HISTORY: Have ever	been treated	for any of the f	following:	
heartlungsliver	kidr	neyulc	ersastl	ıma_
arthritisblood pressure (high	/low)diab	oetes (insulin Y	//N)seiz	zures
strokecancerthyro	id	bleeding di	isorderHIV	V/AIDSHepatitis
depression/anxietyGER	D/REFLUX	hia	tal hernia	
Please describe all yes answers:				
Have you EVER been treated for sleep apn	nea or worn th	e CPAP machi	ine at night? Y/	N
If yes, please call us at 609-581-6230.				
Please list all your daily medications include	ding over the	counter, vitam	ins and supplen	nents:
PAST SURGICAL HISTORY: Please li	st all surgery	(s) you have ha	ad:	

Have you or a family member had any complications from anesthesia? Y/N

SOCIAL HISTORY: Tobacco: No_Yes__ How much? ____ How many years?__ Quit? (when) ____ Alcohol use: heavy moderate social occasional none Recreational drug use? Y/N Do you wear glasses/contacts? Y/N Do you have any dentures/loose teeth/caps/bridges/braces? Permanent Removable Females: Last menstrual period Pregnant? Y/N **INSTRUCTIONS: STOP** for seven days before surgery (unless indicated otherwise by your physician): Aspirin/ Aspirin products Coumadin Plavix Aggrenox **STOP** for three days before surgery (unless indicated otherwise by your physician): NSAIDs Ibuprofen Motrin Celebrex Mobic

The night before surgery:

Naproxen

If you take long-acting insulin in the evening (NPH, Lantus, etc.) only take ½ the dose!!!

Relafen

The morning of surgery:

Do **not** take any insulin or mediation for diabetes, we will check your sugar here.

Take your morning dose of blood pressure, heart, or thyroid medicine with a small sip of water.

Indomethacin Lodine

Nothing to eat or drink after midnight. This includes candy, mints, and gum.

No drinking alcohol, smoking or illicit drug use 24 hours prior to surgery.

Leave all jewelry and valuables at home.

All vitamins/supplements

Naprosyn

Please refrain from wearing perfume/ cologne the morning of surgery.

Bring any equipment as ordered.

Please email, fax or mail this form to:

New Jersey Surgery Center

1225 Whitehorse-Mercerville Rd Fax: 609-585-0527 Building D, Suite 209 Phone: 609-581-6230

Mercerville, NJ 08619 Email: NJSC.PAT@amsurg.com

Attn: Pre-Op Department